

Analysis of Change
*Wetland Surface Elevation Table (SET) Data
for
Great Bay National Estuarine Research Reserve (GRB)
New Hampshire, 2011-2017*



To learn more about project, project team, and related products visit:
nerrssciencencollaborative.org/project/Cressman18

Contact: Kim Cressman, Grand Bay National Estuarine Research Reserve,
kimberly.cressman@dmr.ms.gov

SET Rate Calculations

2020-02-26

This document was designed for use by National Estuarine Research Reserve (NERR) technical staff and may also be useful to other Surface Elevation Table (SET) data analysts. It is meant to calculate, visualize, and describe rates of elevation change at SET stations in a given area (such as an individual NERR or reserve component). These products are output from a 2018-2019 NERRS Science Collaborative Catalyst Grant project nicknamed "SETr". For information on options to customize this report, see the document "Guide to the SETr Workflow" in the main folder of SETr outputs.

Background on the data

This analysis was run on **grbset_processed.csv** on 2020-02-26.

In this report, the user may have chosen to exclude data associated with certain QA/QC codes. Any values that have been removed are listed in the appropriate section below.

Data and metadata setup

We start by reading in the long SET measurement dataset, converted from other formats by earlier SETr scripts. See the "Guide to the SETr Workflow" document for more detail on data formatting. In this step, pin heights were converted to mm if they were previously in cm.

This dataset spans the dates **2011-08-11** to **2017-12-05**.

If custom start and/or end dates were specified for this analysis, the dataset is subsetted here and the chosen date range will be printed below this paragraph. These options can be changed in the file `metadata/user_defined_inputs.xlsx`, general tab.

QA/QC codes

In the same `user_defined_inputs` spreadsheet, certain QA/QC codes may have been specified for removal in this analysis. In the `qaqc_codes` worksheet, they were labeled with “-3”, to be consistent with SWMP’s water quality/weather/nutrient “reject” flagging. If such values are present, they will be printed below, then turned into NAs so they are not used in the analysis.

Note that the R code dealing with these flags looks for an EXACT match in the `qaqc_codes` worksheet, and matches are case-sensitive (all codes should be in all caps). If “LHE” is specified for removal, only values with “LHE” will be removed. “LHE CB” and “CRM LHE” will *not* be removed if only “LHE” is specified; combinations of codes need to be specified on their own line in the input spreadsheet. If you are using a new combination of codes, you may need to create a new row in that spreadsheet See the “Guide to the SETr Workflow” document for more information.

```
## No QA/QC codes were labeled for exclusion. All data values are present.
```

File Matching checks

If any mismatches in SET stations between the data and metadata files are present, they will be noted below. The user of this report will need to make any necessary changes in the data/metadata files.

```
## Warning: The following SET IDs exist in your metadata, but not in your
## data: SFU1, SFU2, SFD1
```

Background information

Reserve-level characteristics

- The local, long-term rate of sea level change is **2.82 +/- 0.16 mm/yr**.
- This rate is reported by Boston, Massachusetts, NWLON station number 8443970 based on data from **1921 to 2017**.
- A shorter-term rate of water level change based on 19 years of data from the same NWLON station, using the same methods NOAA uses to calculate long-term sea level rise (ARIMA 1,0,0), is **4.92 +/- 1.89 mm/yr**. This date range is from **1998 to 2017**.
- The technical report on NOAA’s calculation of long-term SLR trends, *Technical Report NOS CO-OPS 053 - Sea Level Variations of the United States 1854-2006* can be found here (accessed 2020-02-25):
https://tidesandcurrents.noaa.gov/publications/Tech_rpt_53.pdf

SET-level characteristics

Setting

	SET_ID	Type	Main_Veg
	GBF 1	Deep ROD SET	Spartina patens
	GBF 2	Deep ROD SET	Spartina patens
	GBF 3	Deep ROD SET	Spartina patens
	SFD1	Deep ROD SET	Spartina alterniflora
	SFD2	Deep ROD SET	Spartina alterniflora
	SFU1	Deep ROD SET	Spartina alterniflora
	SFU2	Deep ROD SET	Spartina alterniflora
	SPD1	Deep ROD SET	Spartina patens
	SPD2	Deep ROD SET	Spartina alterniflora
	SPSETT5	Deep ROD SET	Spartina patens
	SPU1	Deep ROD SET	Distichilis spicata
	SPU2	Deep ROD SET	Distichilis spicata

Sampling Information


set_id	first_sampled	last_sampled	years_sampled	sample_events
GBF 1	2011-12-05	2017-11-29	5.985	11
GBF 2	2011-12-05	2017-11-29	5.985	11
GBF 3	2011-12-05	2017-11-29	5.985	11
SFD2	2012-05-17	2016-10-05	4.386	5
SPD1	2011-08-11	2017-12-05	6.319	10
SPD2	2014-04-17	2017-12-05	3.636	6
SPSETT5	2012-03-27	2017-11-29	5.676	9
SPU1	2011-08-11	2017-12-05	6.319	10
SPU2	2011-08-11	2017-12-05	6.319	10

Cumulative Change Snapshot

Take a look at your overall change since the first reading - make sure the change looks generally linear, and make sure there are no big breaks in the data that could influence the

outputs. Output will be generated even if it is not appropriate - **it is up to you to use discretion and make sure a linear model is appropriate for the data!**


In the graphics below, the first (baseline) measurement for each pin was subtracted from every subsequent reading, to yield change-since-baseline at a pin level, for all dates. Then pins on an arm were averaged together for each date to yield change-since-baseline at the arm level. Finally, the arms for each date were averaged together to yield change-since-baseline for the SET as a whole. This is a slightly different approach than the rate calculations performed below but gives an almost identical point estimate for rate of change, and makes it easy to put change since baseline on a plot.



The above graph is saved as:

R_output/figures/cumu_change_plots/cumu_change_noLine.png

Graphs for each SET individually are not shown here but have been saved in
R_output/figures/cumu_change_plots/individual_sets



The above graph is saved as:

R_output/figures/cumu_change_plots/cumu_change_withLine.png

Graphs for each SET individually are not shown here but have been saved in
R_output/figures/cumu_change_plots/individual_sets


Plus Sea Level Rise Line

This is an oversimplification of sea level rise: the slope is that of long-term, local SLR, calculated by NOAA COOPS at the NWLON station closest to the reserve. This line does not account for accelerating sea level rise or site-specific processes, and these graphs do not include actual water level data.

These graphs are meant to illustrate and provide context for SET elevation change relative to long-term SLR.

Cumulative Change since first reading

red line has slope of long-term sea level rise; blue line has slope of SET change




The above graph is saved as:

R_output/figures/cumu_change_plots/cumu_change_withLineAndSLR.png

Graphs for each SET individually are not shown here but have been saved in
R_output/figures/cumu_change_plots/individual_sets

Cumulative Change since first reading
red line has the slope of long-term sea level rise




The above graph is saved as:

R_output/figures/cumu_change_plots/cumu_change_withSLR.png

Graphs for each SET individually are not shown here but have been saved in
R_output/figures/cumu_change_plots/individual_sets

Cumulative Change since first reading

light gray represents data; blue has slope of SET change;
red has slope of long-term sea level rise



The above graph is saved as:

R_output/figures/cumu_change_plots/cumu_change_linesOnly.png

Graphs for each SET individually are not shown here but have been saved in
R_output/figures/cumu_change_plots/individual_sets

Cumulative change along the elevation gradient

The following graph shows cumulative elevation change at all SETs in one graph panel. If NAVD88 information was present in metadata, the first reading at each SET was placed at that point on the y-axis. Otherwise, the SET cumulative change lines start at 0.

Available elevation information varied between reserves in this project; **this graph is meant to show changes along a relative elevation gradient only**. We have not overlaid water levels because we could not always be sure of the appropriate offset.

SET cumulative change

Each SET's position on the y-axis was determined by its NAVD88 elevation



The above graph is saved as:

R_output/figures/cumu_change_plots/cumu_change_NAVD88.png

Rate Calculations

From this point on, **only SETs with 5 or more measurements over 4.5 or more years will be analyzed.** If you do not have any SETs that have been measured for this amount of time, you will NOT see analyses or graphs below.

SETs that are included in the following analyses

```
## GBF 1, GBF 2, GBF 3, SPD1, SPSETT5, SPU1, SPU2
```

SETs that are not included in analyses

```
## SFD2, SPD2
```

Statistical methodology

Rates of elevation change at each SET were generated using random-intercept linear mixed models. See Zuur et al. (2009) and Cahoon et al. (2019) for details.

Linear mixed models extend traditional linear regression models by allowing for the inclusion of both fixed and random effects. *These types of models are particularly useful when the data are structured hierarchically*, as with SET data. Data for each SET is analyzed separately using pin height as the response variable; arm and pin (nested in arm) are treated as random effects; and date is considered a numeric covariate. As both the intercept and slope include random effects, methods other than least squares must be employed.

For each SET, we initially considered two LMMs, as in Cahoon et al. (2019): a LMM that includes a random intercept (with a random effects for arm and for pin nested in arm) and a LMM that includes both a random slope and a random intercept (with random effects for arm and for pin nested in arm). For many SETs, we observed that the random intercept model fit better. At other SETs, the random slope and intercept model produced better fitting models (based on AIC). However, the resulting point estimates showed only small differences between the two approaches. As the random intercept models did not require the same degree of scrutiny when model fitting, and did not cause as many script-running errors, we exclusively employed random intercept models in these automated R scripts. For more detailed analyses at a smaller level, we recommend consideration of both models.

General recommendations on analysis of SET data

- Before analysis, the analyst should carefully examine plots of the data to ensure that a linear model is appropriate. *If the points exhibit curvature or if there is some sort of a changepoint, this type of analysis may not be appropriate.* Graphs are provided in this report to help with this determination.
- The analyst should also look for highly influential observations. One way that this could occur is when there is a *large temporal gap* in the data.
- Be aware that, though we have done our best to appropriately describe uncertainty in the following rate calculations, the analyses below do not account for *temporal*

dependence in the data. This could mean that confidence intervals are narrower than they should be and may be something the analyst wishes to address in future analyses.

- The analyst should employ a statistically valid model fitting strategy. When using Linear Mixed Models (LMMs), as we do below, this should also include addressing issues such as:
 - ensuring convergence of the numerical optimization
 - exploring sensitivity to starting values in the optimization procedure
 - determining whether a random slope model or a random slope/random intercept model is most appropriate, and
 - verifying model assumptions.

Statistical details of this analysis

For this analysis, models were fit in R, using the `lme()` function in the `nlme` package (Pinheiro et al. 2019). Confidence intervals were generated using the `intervals()` function, also in the `nlme` package.

Variable names within each SET were:

- **response variable:** pin_height
- **fixed effect:** date
- **random effects:** arm_position, pin_number (with pin_number nested in arm_position)

All calculations generated output in *mm/day* and these rates were converted to *mm/yr* by multiplying by 365.25, to account for leap years.

Calculated rates of elevation change

reserve	set_id	rate	CI_low	CI_high
GRB	GBF 1	2.028	1.698	2.358
GRB	GBF 2	1.553	1.340	1.766
GRB	GBF 3	2.203	2.017	2.390
GRB	SPD1	2.561	2.289	2.833
GRB	SPSETT5	1.305	0.984	1.626
GRB	SPU1	1.162	1.006	1.318
GRB	SPU2	1.254	1.031	1.477

Additional model diagnostics

reserve	set_id	sigma	AIC	BIC	logLik
GRB	GBF 1	6.431	2690.348	2710.230	-1340.174
GRB	GBF 2	4.148	2350.912	2370.794	-1170.456
GRB	GBF 3	3.639	2217.647	2237.529	-1103.824
GRB	SPD1	5.301	2368.922	2388.324	-1179.461
GRB	SPSETT5	5.366	2120.688	2139.560	-1055.344
GRB	SPU1	3.033	2001.356	2020.759	-995.678
GRB	SPU2	4.347	2231.033	2250.436	-1110.517

Comparisons to 0 and SLR

In the following tables and graphics, rates of elevation change at each SET are compared to rates of water level change (SLR = long-term sea level rise; 19yr = water level change over a 19 year period) by investigating whether confidence intervals overlap. This method of comparison was chosen because different methods were used to calculate rates for sea level rise (ARIMA) and SET elevation change (LMMs), using data from different sources. We note that each individual interval has 95% confidence associated with it, and conclusions that are made based on pairwise comparison of these intervals will not necessarily be equivalent to conducting a formal hypothesis test for a difference at the 5% level (Schenker and Gentleman, 2001).

For comparisons to 0, if the SET's 95% CI does not include 0, this *is* equivalent to a formal hypothesis test. For consistency, we are using the same terminology, involving overlapping CIs for all comparisons.

Overall Summary Table

In this table, the SETs are ordered by their rate of change, from lowest at the top to highest at the bottom. The rate and 95% CI (all in mm/yr) for each SET are provided. The last 6 columns show comparisons to: zero (is elevation at the SET changing?), long-term SLR, and 19-year water level change. Each comparison takes up two columns: a comparison of point estimates (is the SET rate higher or lower than what it is compared to), and whether or not the confidence intervals overlap.

SET	Rate	95% CI		Compared to 0		Compared to SLR		Compared to 19-yr change	
		mm/yr	lower	upper	point	CI overlap?	point	CI overlap?	point
SPU1	1.162	1.006	1.318	higher	no	lower	no	lower	no

SET	Rate	95% CI		Compared to 0		Compared to SLR		Compared to 19-yr change	
		mm/yr	lower	upper	point	CI overlap?	point	CI overlap?	point
SPU2	1.254	1.031	1.477	higher	no	lower	no	lower	no
SPSETT5	1.305	0.984	1.626	higher	no	lower	no	lower	no
GBF 2	1.553	1.340	1.766	higher	no	lower	no	lower	no
GBF 1	2.028	1.698	2.358	higher	no	lower	no	lower	no
GBF 3	2.203	2.017	2.390	higher	no	lower	no	lower	no
SPD1	2.561	2.289	2.833	higher	no	lower	yes	lower	no

Increasing/Decreasing (Comparison to 0)

The following tables break the SETs into groups where the rate of SET elevation change is *lower than / higher than / not different from 0*. *Lower than* and *higher than* tables imply that the 95% confidence intervals for the SET's rate of elevation change do not include 0. *Not different from* means that 0 is included.

SET Elevation Change < 0 mm/yr

reserve	set_id	rate	CI_low	CI_high
---------	--------	------	--------	---------

SET Elevation Change > 0 mm/yr

reserve	set_id	rate	CI_low	CI_high
GRB	GBF 1	2.028	1.698	2.358
GRB	GBF 2	1.553	1.340	1.766
GRB	GBF 3	2.203	2.017	2.390
GRB	SPD1	2.561	2.289	2.833
GRB	SPSETT5	1.305	0.984	1.626
GRB	SPU1	1.162	1.006	1.318
GRB	SPU2	1.254	1.031	1.477

SET Elevation Change 95% CI Includes 0 mm/yr

reserve	set_id	rate	CI_low	CI_high
---------	--------	------	--------	---------

Sea Level Rise Comparisons

Period of Record (long-term SLR)

The long-term local rate of sea level rise is **2.82 +/- 0.16 mm/yr**.

This rate is reported by Boston, Massachusetts, NWLON station number 8443970 based on data from 1921 to 2017.

The following tables break the SETs into groups where the rate of SET elevation change is *lower than / higher than / not different from* this SLR rate. *Lower than* and *higher than* tables imply that 95% confidence intervals do not overlap between the SET and SLR. *Not different from* means that confidence intervals *do* overlap.

This method of comparison was chosen because different methods were used to calculate rates for sea level rise (ARIMA) and SET elevation change (LMMs) using data from different sources. We note that each individual interval has 95% confidence associated with it, and conclusions that are made based on pairwise comparison of these intervals will not necessarily be equivalent to conducting a formal hypothesis test for a difference at the 5% level (Schenker and Gentleman, 2001).

SET Elevation Change < SLR; CIs don't overlap

reserve	set_id	rate	CI_low	CI_high
GRB	GBF 1	2.028	1.698	2.358
GRB	GBF 2	1.553	1.340	1.766
GRB	GBF 3	2.203	2.017	2.390
GRB	SPSETT5	1.305	0.984	1.626
GRB	SPU1	1.162	1.006	1.318
GRB	SPU2	1.254	1.031	1.477

SET Elevation Change > SLR; CIs don't overlap

reserve	set_id	rate	CI_low	CI_high
---------	--------	------	--------	---------

SET Elevation Change and SLR CIs overlap

reserve	set_id	rate	CI_low	CI_high
GRB	SPD1	2.561	2.289	2.833

19-year water level change

The local, 19-year rate of water level change is **4.92 +/- 1.89 mm/yr**.

This rate uses data reported by Boston, Massachusetts, NWLON station number 8443970 based on data from 1998 to 2017.

The following tables break the SETs into groups where the rate of SET elevation change is *lower than / higher than / not different from* this 19-year rate. *Lower than* and *higher than* tables imply that 95% confidence intervals do not overlap between the SET and water level change. *Not different from* means that confidence intervals *do* overlap.

This method of comparison was chosen because different methods were used to calculate rates for sea level rise (ARIMA) and SET elevation change (LMMs) using data from different sources. We note that each individual interval has 95% confidence associated with it, and conclusions that are made based on pairwise comparison of these intervals will not necessarily be equivalent to conducting a formal hypothesis test for a difference at the 5% level (Schenker and Gentleman, 2001).

SET Elevation Change < 19-year water level change; CIs don't overlap

reserve	set_id	rate	CI_low	CI_high
GRB	GBF 1	2.028	1.698	2.358
GRB	GBF 2	1.553	1.340	1.766
GRB	GBF 3	2.203	2.017	2.390
GRB	SPD1	2.561	2.289	2.833

reserve	set_id	rate	CI_low	CI_high
GRB	SPSETT5	1.305	0.984	1.626
GRB	SPU1	1.162	1.006	1.318
GRB	SPU2	1.254	1.031	1.477

SET Elevation Change > 19-year water level change; CIs don't overlap

reserve	set_id	rate	CI_low	CI_high
---------	--------	------	--------	---------


SET Elevation Change and 19-year water level change CIs overlap

reserve	set_id	rate	CI_low	CI_high
---------	--------	------	--------	---------

Graphical Comparisons to Sea Level Rise and 0


If dominant vegetation was provided in the metadata document, the following graphs will be provided both with and without points colored by vegetation type. If any vegetation entries were missing, the vegetation point-coloration will NOT be provided. You can generate them by adding vegetation to the “CoDominant Species 1” column of the --- `set_metadata.xlsx` document.

Sites in alphabetical order, R's default



```
## The above graph is saved as: R_output/figures/summary_plots/summary_plot.png
```

```
## The above graph is saved as: R_output/figures/summary_plots/summary_plot_veg.png
```




```
## The above graph is saved as: R_output/figures/summary_plots/summary_plot_
bothSLRs.png


## The above graph is saved as: R_output/figures/summary_plots/summary_plot_
veg_19yr.png
```

Ordered (categorically) by NAVD88 elevation

If NAVD88 elevations were provided in the metadata, two more versions of the graph above are produced below. The SETs are ordered along the y-axis from highest to lowest elevation.



```
## The above graph is saved as: R_output/figures/summary_plots/summary_plot_
bothSLRs_navd88ordering.png
```



```
## The above graph is saved as: R_output/figures/summary_plots/summary_plot_
bothSLRs_navd88ordering_veg.png
```


Sites by a user-specified order

If the `numerical_order` column of the metadata was filled out, the order of SETs in the graphics below will be in that order, as well as labeled with user-friendly names. Note that no plot will be produced if there are any NAs (missing values) in the metadata fields `numerical_order` or `user_friendly_set_name`.

MAPS

The way the maps are rendered below (and even *if* they are rendered) depends on several things specific to your computer. The script `R_scripts/04_interact_maps.R` will let you interact with the maps, and you can either take a screenshot or use the Export command from RStudio's Viewer pane to save a version that looks better. Even if no output was generated in this Word document, you should still be able to use the interactive script to generate maps.


Comparisons to 0



Comparisons to long-term SLR



Comparisons to 19-yr water level change



References

- Cahoon, D.R., J.C. Lynch, C.T. Roman, J.P. Schmit, and D.E. Skidds. 2019. Evaluating the Relationship Among Wetland Vertical Development, Elevation Capital, Sea-Level Rise, and Tidal Marsh Sustainability. *Estuaries and Coasts* 42:1-15. <https://doi.org/10.1007/s12237-018-0448-x>
- Pinheiro J, Bates D, DebRoy S, Sarkar D, R Core Team. 2019. *nlme: Linear and Nonlinear Mixed Effects Models*. R package version 3.1-140, <https://CRAN.R-project.org/package=nlme>
- Schenker, N. and J.F. Gentleman. 2001. On Judging the Significance of Differences by Examining the Overlap Between Confidence Intervals. *The American Statistician* 55(3):182-186. <https://doi.org/10.1198/000313001317097960>
- Zuur, A.F., E.N. Ieno, N.J. Walker, A.A. Saveliev, and G.M. Smith. 2009. Mixed effects models and extensions in ecology with R. New York: Springer.